

NOAA Technical Report NMFS SSRF-699



**Seasonal Surface Currents Off the
Coasts of Vancouver Island and
Washington as Shown by
Drift Bottle Experiments, 1964-65**

W. JAMES INGRAHAM, JR. and JAMES R. HASTINGS

SEATTLE, WA
MAY 1976

noaa

NATIONAL OCEANIC AND
ATMOSPHERIC ADMINISTRATION

National Marine
Fisheries Service

NOAA TECHNICAL REPORTS

National Marine Fisheries Service, Special Scientific Report—Fisheries Series

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for optimum use of the resources. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The Special Scientific Report—Fisheries series was established in 1949. The series carries reports on scientific investigations that document long-term continuing programs of NMFS, or intensive scientific reports on studies of restricted scope. The reports may deal with applied fishery problems. The series is also used as a medium for the publication of bibliographies of a specialized scientific nature.

NOAA Technical Reports NMFS SSRF are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained (unless otherwise noted) from D83, Technical Information Division, Environmental Science Information Center, NOAA, Washington, D.C. 20235. Recent SSRF's are:

619. Macrozooplankton and small nekton in the coastal waters off Vancouver Island (Canada) and Washington, spring and fall of 1963. By Donald S. Day. January 1971, iii + 94 p., 19 figs., 13 tables.
620. The Trade Wind Zone Oceanography Pilot Study. Part IX: The sea-level wind field and wind stress values, July 1963 to June 1965. By Gunter R. Seckel. June 1970, iii + 66 p., 5 figs.
621. Predation by sculpins on fall chinook salmon, *Oncorhynchus tshawytscha*, fry of hatchery origin. By Benjamin G. Patten. February 1971, iii + 14 p., 6 figs., 9 tables.
622. Number and lengths, by season, of fishes caught with an otter trawl near Woods Hole, Massachusetts, September 1961 to December 1962. By F. E. Lux and F. E. Nichy. February 1971, iii + 15 p., 3 figs., 19 tables.
623. Apparent abundance, distribution, and migrations of albacore, *Thunnus alalunga*, on the North Pacific longline grounds. By Brian J. Rothschild and Marian Y. Y. Yong. September 1970, v + 37 p., 19 figs., 5 tables.
624. Influence of mechanical processing on the quality and yield of bay scallop meats. By N. B. Webb and F. B. Thomas. April 1971, iii + 11 p., 9 figs., 3 tables.
625. Distribution of salmon and related oceanographic features in the North Pacific Ocean, spring 1968. By Robert R. French, Richard G. Bakkala, Masanzo Osako, and Jun Ito. March 1971, iii + 22 p., 19 figs., 3 tables.
626. Commercial fishery and biology of the freshwater shrimp, *Macrobrachium*, in the Lower St. Paul River, Liberia, 1952-53. By George C. Miller. February 1971, iii + 13 p., 8 figs., 7 tables.
627. Calico scallops of the Southeastern United States, 1959-69. By Robert Cummins, Jr. June 1971, iii + 22 p., 23 figs., 3 tables.
628. Fur Seal Investigations, 1969. By NMFS, Marine Mammal Biological Laboratory. August 1971, 82 p., 20 figs., 44 tables, 23 appendix A tables, 10 appendix B tables.
629. Analysis of the operations of seven Hawaiian skipjack tuna fishing vessels, June-August 1967. By Richard N. Uchida and Ray F. Sumida. March 1971, v + 25 p., 14 figs., 21 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
630. Blue crab meat. I. Preservation by freezing. July 1971, iii + 13 p., 5 figs., 2 tables. II. Effect of chemical treatments on acceptability. By Jurgen H. Strasser, Jean S. Lennon, and Frederick J. King. July 1971, iii + 12 p., 1 fig., 9 tables.
631. Occurrence of thiaminase in some common aquatic animals of the United States and Canada. By R. A. Greig and R. H. Gnaedinger. July 1971, iii + 7 p., 2 tables.
632. An annotated bibliography of attempts to rear the larvae of marine fishes in the laboratory. By Robert C. May. August 1971, iii + 24 p., 1 appendix I table, 1 appendix II table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
633. Blueing of processed crab meat. II Identification of some factors involved in the blue discoloration of canned crab meat *Callinectes sapidus*. By Melvin E. Waters. May 1971, iii + 7 p., 1 fig., 3 tables.
634. Age composition, weight, length, and sex of herring, *Clupea pallasii*, used for reduction in Alaska, 1929-66. By Gerald M. Reid. July 1971, iii + 25 p., 4 figs., 18 tables.
635. A bibliography of the blackfin tuna, *Thunnus atlanticus* (Lesson). By Grant L. Beardsley and David C. Simmons. August 1971, 10 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
636. Oil pollution on Wake Island from the tanker *R. C. Stoner*. By Reginald M. Gooding. May 1971, iii + 12 p., 8 figs., 2 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
637. Occurrence of larval, juvenile, and mature crabs in the vicinity of Beaufort Inlet, North Carolina. By Donnie L. Dudley and Mayo H. Judy. August 1971, iii + 10 p., 1 fig., 5 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
638. Length-weight relations of haddock from commercial landings in New England, 1931-55. By Bradford E. Brown and Richard C. Hennemuth. August 1971, v + 13 p., 16 figs., 6 tables, 10 appendix A tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
639. A hydrographic survey of the Galveston Bay system, Texas 1963-66. By E. J. Pullen, W. L. Trent, and G. B. Adams. October 1971, v + 13 p., 15 figs., 12 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
640. Annotated bibliography on the fishing industry and biology of the blue crab, *Callinectes sapidus*. By Marlin E. Tagatz and Ann Bowman Hall. August 1971, 94 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
641. Use of threadfin shad, *Dorosoma petenense*, as live bait during experimental pole-and-line fishing for skipjack tuna, *Katsuwonus pelamis*, in Hawaii. By Robert T. B. Iversen. August 1971, iii + 10 p., 3 figs., 7 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
642. Atlantic menhaden *Brevoortia tyrannus* resource and fishery—analysis of decline. By Kenneth A. Henry. August 1971, v + 32 p., 40 figs., 6 appendix figs., 3 tables, 2 appendix tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
643. Surface winds of the southeastern tropical Atlantic Ocean. By John M. Steigler and Merton C. Ingham. October 1971, iii + 20 p., 17 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
644. Inhibition of flesh browning and skin color fading in frozen fillets of yelloweye snapper (*Lutjanus triviro*). By Harold C. Thompson, Jr., and Mary H. Thompson. February 1972, iii + 6 p., 3 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
645. Traveling screen for removal of debris from rivers. By Daniel W. Bates, Ernest W. Murphey, and Martin G. Beam. October 1971, iii + 6 p., 6 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington D.C. 20402.
646. Dissolved nitrogen concentrations in the Columbia and Snake Rivers in 1970 and their effect on chinook salmon and steelhead trout. By Wesley J. Ebel. August 1971, iii + 7 p., 2 figs., 6 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington D.C. 20402.
647. Revised annotated list of parasites from sea mammals caught off the west coast of North America. By L. Margolis and M. D. Dailey. March 1972, iii + 23 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

Continued on inside back cover.

NOAA Technical Report NMFS SSRF-699

Seasonal Surface Currents Off the Coasts of Vancouver Island and Washington as Shown by Drift Bottle Experiments, 1964-65

W. JAMES INGRAHAM, JR. and JAMES R. HASTINGS

SEATTLE, WA
MAY 1976

UNITED STATES
DEPARTMENT OF COMMERCE
Elliot L. Richardson, Secretary

NATIONAL OCEANIC AND
ATMOSPHERIC ADMINISTRATION
Robert M. White, Administrator

National Marine
Fisheries Service
Robert W. Schoning, Director



9 pp

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

	Page
Introduction	1
April 1964	1
July 1964	2
November 1964	4
January 1965	4
Summary	6
Acknowledgments	7
Literature cited	7

Figures

1. Locations of drift bottle releases and recoveries, RV <i>George B. Kelez</i> , 20-28 April 1964	2
2. Locations of drift bottle releases and recoveries, RV <i>George B. Kelez</i> , 20-26 July 1964	2
3. Locations of drift bottle releases and recoveries, RV <i>George B. Kelez</i> , 6-16 November 1964	6
4. Locations of drift bottle releases and recoveries, RV <i>George B. Kelez</i> , 13-17 January 1965	6

Tables

1. Release and recovery locations for drift bottles released in April 1964	3
2. Release and recovery locations for drift bottles released in July 1964	5
3. Release and recovery locations for drift bottles released in November 1964	7
4. Release and recovery locations for drift bottles released in January 1965	8

Seasonal Surface Currents Off the Coasts of Vancouver Island and Washington as Shown by Drift Bottle Experiments, 1964-65

W. JAMES INGRAHAM, JR. and JAMES R. HASTINGS¹

ABSTRACT

Release of a total of 1,044 drift bottles during four periods from April 1964 to January 1965 off the coasts of Washington and Vancouver Island, British Columbia, indicate a seasonal reversal of flow, southward in April and July and northward in November and January, within 200 km (108 nautical miles) of the coast. This study supplements those conducted off the Oregon and California coasts by other agencies.

INTRODUCTION

The Northwest Fisheries Center of the National Marine Fisheries Service, Seattle, Wash. (Biological Laboratory of the Bureau of Commercial Fisheries prior to 1970), has conducted far-ranging research on fisheries problems throughout the subarctic Pacific region from local coastal waters of Washington to the central North Pacific Ocean and the Bering Sea. Oceanographic conditions that affect the distribution and abundance of fish stocks have been studied in conjunction with fisheries research since 1955. Studies of local ocean currents off the coasts of Washington and Vancouver Island were emphasized in the oceanographic research program from 1963 to 1964 (Ingraham 1967). The purpose here is to summarize results of drift bottle experiments in 1964 and 1965 which were not reported by Favorite and Fisk (1971) nor by Fisk (1971).

Although the general oceanic flow may appear as a simple onshore movement, eddies form complex flow patterns within about 500 km (270 nautical miles) of shore as the land boundary causes the current to diverge toward the north and south. These complexities were shown by dynamic computations and from records of water properties for several isolated studies at different times (Doe 1955; Bennett 1959; Reid 1960; Budinger et al. 1964). Other studies, using drift bottles to estimate surface water drift, have shown some details of flow within 200 km (108 nautical miles) of shore where reversals occur, with flow from the north in summer and from the south in winter (Dodimead and Hollister 1958; Schwartzlose 1963; Burt and Wyatt 1964).

Four oceanographic cruises were conducted aboard the RV *George B. Kelez* in April, July, and November 1964 and January 1965, which helped provide areal continuity in nearshore ocean sampling between the Oregon and Canadian coasts.² During these cruises 1,044 drift bottles were released along two lines normal to shore—one from Willapa Bay, Wash., to Cobb Seamount and the other from Cobb

Seamount to Estevan Point, Vancouver Island, B.C. Twelve corked, unballasted, short-necked (17 cm long with a 4½-cm base) drift bottles were released at 21 stations during each cruise. Ballast sand was not placed in the bottles to save handling time. These particular empty bottles already floated about 80% submerged and any improvement would be minor compared to other inherent errors in the method. Only minimum speeds are inferred by drift bottle data because there is no way of knowing how long the bottle was on the beach before recovery. Each contained a folded data card with instructions in English, Russian, and Japanese and four-digit serial numbers ranging from 6001 to 7044. A reward (\$1.00) was offered for the return of each card. Recoveries from each cruise are discussed separately below.

APRIL 1964

From 20 to 28 April 1964, a total of 288 drift bottles was released at 21 stations and 77 recoveries (27%) have been reported (Fig. 1, Table 1). The recoveries suggest two distinct regions with different flow features: 1) a southerly drift at speeds of 15 to 30 cm/s about 167 km (90 nautical miles) wide nearshore and 2) an offshore region of questionable flow from which no recoveries have been reported. These features are the same along both of the tracklines.

The 41 recoveries from the six inshore stations along the trackline normal to the Washington coast were from the Washington and Oregon coasts southeast of the release locations. All 7 of the recoveries from the nearest inshore station were reported from the southern Washington coast, whereas, all of the remaining 34 recoveries were from the Oregon coast. These reflect a dominant southerly flow of surface waters at this time. A northward flow was reported in April 1959 and 1960 (Schwartzlose 1963), and mixed flow toward the north and south was reported in April 1961 and 1962 (Burt and Wyatt 1964).

Along the trackline normal to the west coast of Vancouver Island, 36 recoveries came from 72 releases at the six inshore stations. Of the 15 recoveries reported from the three stations shoreward of the 100-fathom curve, four were recovered on the west coast of Vancouver Island north of the drop location, indicating some localized northerly flow of surface waters along the west coast of Vancouver Island. The remainder of the recoveries from these three stations and 20

¹Northwest Fisheries Center, National Marine Fisheries Service, NOAA, 2725 Montlake Boulevard East, Seattle, WA 98112.

²Ingraham, W. J., Jr., and D. M. Fisk. 1966. Oceanographic observations off the coasts of Washington and British Columbia—April, July, and November, 1964 and January 1965. Unpubl. manuscr., 96 p., Northwest Fisheries Center, Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98112.




Figure 1.—Locations of drift bottle releases and recoveries, RV *George B. Kelez*, 20-28 April 1964.

of the 21 recoveries from the three seaward stations were recovered along the Washington and Oregon coasts. No recoveries to date have been reported from the two offshore stations along this trackline.

On 14 May 1964, a transponding telemetry buoy was launched at lat. 48°00'N, long. 131°53'W (Favorite et al. 1965). An additional 48 drift bottles were released at this site, but no recoveries have been reported. The buoy drifted east-southeast (115° true direction) for 595 km (320 nautical miles) at an average speed of 11 cm/s. This suggests the drift bottles released offshore may have moved onshore but, before coming ashore, they were carried offshore into the eastern edge of the subtropical North Pacific gyre.

JULY 1964

There were 37 recoveries (16%) from 228 releases at the 19 stations occupied during the July 1964 cruise (Fig. 2, Table 2). They numbered less than one-half the recoveries reported for the April 1964 cruise. In general, the fewer returns are consistent with the expected usual July conditions of northerly winds which transport surface water offshore. Recoveries are probably associated with sporadic short-term events in which southerly winds transport bottles that have remained nearshore onto the beach. Features of surface drift are very similar to those of April—a slower southeasterly




Figure 2.—Locations of drift bottle releases and recoveries, RV *George B. Kelez*, 20-26 July 1964.

flow at 5 to 11 cm/s (minimum) within about 185 km (100 nautical miles) of shore and no returns from offshore.

Recoveries were made from only 4 of the 11 stations along the trackline normal to the Washington coast during this cruise. Of the 13 recoveries, 6 drifted from the station closest to shore onto the northern Oregon coast. No recoveries to date have been reported from releases at the next three seaward stations, but four recoveries, ranging from southern Washington to northern Oregon, have been made from the fifth station from shore. From the remaining four stations on this trackline, there were only three recoveries; all were from south of the mouth of the Columbia River, with the recovery from the most seaward station coming from northern California (lat. 38°N).

Almost twice as many recoveries were reported from the trackline normal to the west coast of Vancouver Island as the trackline normal to the Washington coast—24 recoveries from 96 released. Nine recoveries were released from the three stations nearest Vancouver Island, five from the west coast of Vancouver Island. Two of these five recoveries were from northwest of the launch site, indicating some deviation from the general southerly flow along the coast during this time of year. The other three recoveries were made near Estevan Point on Vancouver Island. The 19 other recoveries were made along the coast from central Washington to northern California. It appears that during this season the

Table 1. — Release and recovery locations for drift bottles released in April 1964.

Serial number	Date released	Location		Number released	Number recovered	Serial number	Date recovered	Location	
		Lat. N	Loog. W					Lat. N	Long. W
6001-6012	4/20/64	46°45'	124°20'	12	7	6001	3/10/66	46°26'	124°03'
						6002	4/22/64	46°24'	124°03'
						6003	4/22/64	46°26'	124°03'
						6005	5/1/64	46°18'	124°02'
						6006	4/22/64	46°24'	124°03'
						6009	4/22/64	46°18'	124°02'
						6011	5/7/64	46°19'	124°02'
6013-6024	4/20/64	46°45'	124°35'	12	10	6013	4/26/64	45°35'	123°56'
						6014	4/25/64	45°33'	123°57'
						6015	4/26/64	45°37'	123°53'
						6017	4/26/64	45°28'	123°53'
						6018	4/25/64	45°33'	123°57'
						6019	4/24/64	45°39'	123°56'
						6020	4/24/64	45°39'	123°56'
						6022	4/24/64	45°39'	123°56'
						6023	4/25/64	45°39'	123°56'
						6024	4/25/64	—	— *
						6025-6036	4/20/64	46°45'	124°48'
6028	5/4/64	44°23'	124°03'						
6035	5/8/64	43°57'	124°07'						
6037-6048	4/21/64	46°45'	125°20'	12	9	6037	5/8/64	44°55'	124°00'
						6039	5/8/64	45°10'	123°58'
						6040	5/7/64	44°55'	124°00'
						6041	5/8/64	44°55'	124°00'
						6042	5/7/64	44°55'	124°00'
						6043	5/7/64	44°55'	124°00'
						6045	5/4/64	45°49'	123°57'
						6046	5/8/64	44°55'	124°00'
						6047	5/7/64	45°49'	123°57'
6049-6060	4/21/64	46°45'	125°10'	12	9	6050	5/9/64	44°35'	124°03'
						6051	5/9/64	44°55'	124°00'
						6052	5/5/64	45°39'	123°56'
						6053	7/21/64	43°04'	124°35'
						6054	5/7/64	44°55'	124°00'
						6055	5/6/64	45°45'	123°57'
						6058	5/17/64	42°46'	124°29'
						6059	5/22/64	42°47'	124°29'
						6060	5/3/64	45°49'	123°57'
						6061-6072	4/23/64	46°45'	126°00'
6070	6/9/64	43°57'	124°07'						
5072	12/24/66	—	— *						
6073-6084	4/23/64	46°45'	126°33'	12	0	—	—	—	—
6085-6096	4/25/64	46°45'	127°12'	12	0	—	—	—	—
6097-6108	4/25/64	46°45'	127°48'	12	0	—	—	—	—
6109-6120	4/25/64	46°45'	128°24'	12	0	—	—	—	—
6121-6132	4/26/64	46°45'	129°36'	12	0	—	—	—	—
6133-6144	4/26/64	46°45'	130°34'	12	0	—	—	—	—
6145-6156	4/26/64	47°20'	129°59'	12	0	—	—	—	—
6157-6168	4/27/64	47°53'	129°02'	12	0	—	—	—	—
6169-6180	4/27/64	48°26'	128°07'	12	3	6171	5/24/64	46°14'	124°04'
						6172	5/24/64	46°13'	124°02'
						6174	6/28/64	43°22'	124°15'

Table 1.—Release and recovery locations for drift bottles released in April 1964, continued.

Serial number	Date released	Location		Number released	Number recovered	Serial number	Date recovered	Location	
		Lat. N	Long. W					Lat. N	Long. W
6181-6192	4/27/64	48°58'	127°15'	12	8	6183	5/25/64	47°02'	124°08'
						6184	5/31/64	46°58'	124°00'
						6185	6/26/64	49°09'	125°59'
						6186	7/2/64	47°10'	124°10'
						6187	5/23/64	—	— *
						6188	5/25/64	48°09'	124°44'
						6189	6/13/64	47°52'	124°38'
6193-6204	4/27/64	49°04'	127°07'	12	10	6193	5/23/64	47°41'	124°29'
						6194	5/23/64	47°34'	124°21'
						6195	5/23/64	47°34'	124°21'
						6197	5/28/64	46°47'	124°04'
						6198	5/23/64	47°05'	124°09'
						6199	5/28/64	47°31'	124°20'
						6200	5/28/64	46°31'	124°04'
						6201	5/23/64	47°34'	124°21'
						6202	6/18/64	44°27'	124°04'
						6204	5/23/64	47°34'	124°21'
						6205-6216	4/28/64	49°11'	126°54'
6207	6/20/64	47°41'	124°29'						
6208	10/5/64	50°48'	128°26'						
6209	5/30/64	47°57'	129°40'						
6210	5/27/64	48°09'	124°44'						
6211	6/21/64	46°26'	124°03'						
6213	5/28/64	47°05'	124°09'						
6214	6/6/64	48°22'	124°37'						
6216	6/18/64	47°33'	124°21'						
6217-6228	4/28/64	49°16'	126°46'	12	5				
						6220	5/12/64	50°08'	127°55'
						6223	6/1/64	46°47'	124°04'
						6227	7/26/64	50°06'	127°51'
						6228	5/29/64	47°04'	124°08'
6229-6240	4/28/64	49°20'	126°37'	12	1	6229	9/27/65	49°37'	126°46'
6241-6288	5/14/64	48°00'	131°53'	48	0	—	—	—	—

*Card returned—unable to ascertain recovery location.

farther from shore the releases were made, the farther southward the travel before recovery.

NOVEMBER 1964

The cruise during November 1964 yielded the fewest recoveries (9%), 24 of the 276 released (Fig. 3, Table 3). The few recoveries prevent defining any clear pattern of circulation for this time of the year, but a generally slow northward flow is evident from all but one recovery.

There were 18 recoveries (12%) from 156 releases along the trackline normal to the Washington coast. Two of the bottles released at the station closest to shore drifted into the Strait of Juan de Fuca and were recovered at the southern tip of Vancouver Island. These recoveries were reported 5 wk after their release and suggest a general northerly flow along the coast at 5 to 8 cm/s (minimum) with an appreciable onshore component, particularly at the entrance to the Strait of Juan de Fuca. The single recovery from the adjacent station along this trackline was reported from Montague Island in Prince William Sound, Alaska (lat. 59°50'N, long. 147°45'W), whereas the recovery from the station nearest

Cobb Seamount was reported from the central portion of the west coast of Vancouver Island. Of the 10 remaining stations along this trackline, recoveries have been reported from only 3, and all but 1 of the 14 recoveries were from the Washington or Vancouver Island coasts.

Only six recoveries (5%) have been reported from 120 releases at the 10 stations along the trackline normal to Vancouver Island. All of these were from only five stations, and all recoveries were made north of the release area; four of the six recoveries were in Laredo Sound (about lat. 52°30'N, long. 129°W).

JANUARY 1965

The winter cruise from 13 to 18 January 1965 was unique because recoveries were made from nearly all the 21 release stations. A total of 252 drift bottles was released and 76 were recovered (Fig. 4, Table 4)—a 30% recovery. Speed and direction of surface flow are indicated by first recoveries from each station that fell into three different time groups: 1) recoveries within 11 to 24 days released within 130 km (70 nautical miles) offshore indicate the extension of the

Table 2. — Release and recovery locations for drift bottles released in July 1964.

Serial number	Date released	Location		Number released	Number recovered	Serial number	Date recovered	Location	
		Lat. N	Long. W					Lat. N	Long. W
6289-6300	7/20/64	46°45'	124°18'	12	6	6289	8/22/64	45°26'	123°58'
						6291	8/21/64	45°24'	123°58'
						6293	8/8/64	45°12'	123°56'
						6295	8/9/64	45°55'	123°53'
						6299	8/13/64	45°28'	123°56'
						6300	8/12/64	45°03'	123°59'
6301-6312	7/20/64	46°45'	124°36'	12	0	—	—	—	—
6313-6324	7/21/64	46°45'	124°47'	12	0	—	—	—	—
6325-6336	7/21/64	46°45'	125°10'	12	0	—	—	—	—
6337-6348	7/21/64	46°45'	125°20'	12	4	6339	8/9/64	46°42'	124°05'
						6341	8/13/64	46°25'	124°03'
						6347	8/15/64	45°40'	123°56'
						6348	8/ /64 ¹	46°09'	124°00'
6349-6360	7/22/64	46°45'	126°00'	12	0	—	—	—	—
6361-6372	7/22/64	46°45'	126°33'	12	2	6361	12/24/64	46°02'	123°56'
						6367	12/11/64	44°27'	124°04'
6373-6384	7/22/64	46°45'	127°11'	12	0	—	—	—	—
6385-6396	7/23/64	46°45'	128°24'	12	1	6388	1/1/65	38°00'	123°00'
6397-6408	7/23/64	46°45'	129°36'	12	0	—	—	—	—
6409-6420	7/23/64	46°45'	130°33'	12	0	—	—	—	—
6421-6432	7/24/64	47°20'	129°57'	12	0	—	—	—	—
6433-6444	7/24/64	47°53'	129°15'	12	3	6435	12/3/64	40°43'	124°14'
						6436	1/28/65	38°04'	123°00'
						6438	1/11/69	43°50'	124°08'
6445-6456	7/25/64	48°27'	128°09'	12	4	6448	12/31/64	43°51'	124°09'
						6449	11/15/64	41°23'	124°03'
						6451	11/28/64	43°20'	124°19'
						6455	12/29/64	44°54'	124°01'
						—	—	—	—
6457-6468	7/25/64	49°20'	126°37'	12	1	6463	8/24/64	50°08'	127°55'
6469-6480	7/26/64	49°14'	126°46'	12	1	6476	8/23/64	50°12'	127°49'
6481-6492	7/26/64	49°10'	126°56'	12	7	6481	4/3/66	49°25'	126°35'
						6482	11/2/64	49°15'	126°10'
						6483	10/20/64	49°25'	126°23'
						6487	12/1/64	45°25'	123°57'
						6490	12/31/64	46°26'	124°03'
						6491	9/25/64	46°52'	124°07'
						6492	11/10/64	43°18'	124°23'
						—	—	—	—
6493-6504	7/26/64	49°06'	127°00'	12	5	6494	11/23/68	45°18'	123°59'
						6495	12/30/64	46°30'	124°02'
						6497	9/20/64	45°53'	123°59'
						6501	9/20/64	45°21'	123°57'
						6502	10/14/64	43°50'	124°08'
6505-6516	7/26/64	49°00'	127°10'	12	3	6508	10/17/64	43°22'	124°17'
						6509	12/21/64	45°16'	124°00'
						6514	12/1/64	43°56'	124°08'

¹ Card returned, without date, during August 1964.




Figure 3.—Locations of drift bottle releases and recoveries, RV *George B. Kelez*, 6-16 November 1964.




Figure 4.—Locations of drift bottle releases and recoveries, RV *George B. Kelez*, 13-17 January 1965.

Davidson Current flowing northward off the coast of Washington at 11 to 15 cm/s (minimum) and flowing toward the northwest off the coast of Vancouver island at 13 to 17 cm/s (minimum); 2) recoveries 41 to 44 days after release indicate the current still had a net northerly component as far as 445 km (240 nautical miles) offshore; and 3) recoveries 60 to 70 days after release near Cobb Seamount were reported both north and south of their release latitude. Many late recoveries (70 days) were made from most of the stations beyond 130 km (70 nautical miles), which showed a wide range of recovery points from northern Washington to southern Oregon. Their significance is questionable, but multiple late recoveries probably suggest a large dispersion of bottles from any one release station with a possible long residence close to shore once the nearshore zone is reached, followed by sporadic recoveries depending on the local onshore transport.

SUMMARY

As part of an ocean survey of local seasonal conditions off the coasts of Washington and Vancouver Island, 12 drift bottles were released at each of 21 stations along two lines from Cobb Seamount normal to the shorelines of each coast during April, July, and November 1964 and January 1965. These experiments extend our knowledge of drift of coastal

surface water northward of the area of earlier studies by Scripps Institute of Oceanography and Oregon State University, providing data for another time period.

Early recoveries from both lines of stations within about 60 days of release clearly showed the dominant seasonal patterns of flow nearshore. Of the April releases 27% were recovered, compared to lower recoveries in similar studies off Oregon (13%) and off California (5%). The percentage of recoveries decreased during July to 16% and was lowest in November at 9%, but January recoveries were highest at 30%. Within about 150 to 200 km (81 to 108 nautical miles) of shore, the surface drift was southerly during April and July but speeds of first returns decreased from a maximum of 15 to 30 cm/s (minimum) in April to 5 to 11 cm/s (minimum) in July. The direction of flow reversed sometime before mid-autumn, for a well-developed northward flow at speeds of 5 to 8 cm/s (minimum) extended along both the Washington and Vancouver Island coasts beyond lat. 50°N in November. The northward flow was also present in January but speeds had doubled. Very little may be inferred about nearshore flow from bottles released beyond 200 km (108 nautical miles) from the coast. Those released in November or January may be recovered but move in a long complex path. Those released in April or July appear to have been caught in the North Pacific gyre as shown by Dodimead and Hollister (1958).

Table 3. — Release and recovery locations for drift bottles released in November 1964.

Serial number	Date released	Location		Number released	Number recovered	Serial number	Date recovered	Location	
		Lat. N	Long. W					Lat. N	Long. W
6517-6528	11/6/64	46°45'	124°19'	12	2	6517	12/14/64	48°25'	123°24'
						6518	12/12/64	48°23'	123°55'
6529-6540	11/7/64	46°45'	124°36'	12	1	6534	12/4/64	59°50'	147°45'
6541-6552	11/7/64	46°45'	124°48'	12	1	6545	12/2/68	48°57'	125°35'
6553-6564	11/8/64	46°45'	124°59'	12	0	—	—	—	—
6565-6576	11/8/64	46°45'	125°10'	12	0	—	—	—	—
6577-6588	11/9/64	46°45'	125°20'	12	0	—	—	—	—
6589-6600	11/9/64	46°45'	126°00'	12	4	6591	12/12/64	47°54'	124°37'
						6593	12/12/64	48°00'	124°41'
						6595	12/12/64	48°00'	124°41'
						6599	12/12/64	47°54'	124°37'
6601-6612	11/10/64	46°45'	126°33'	12	0	—	—	—	—
6613-6624	11/10/64	46°45'	127°11'	12	0	—	—	—	—
6625-6636	11/10/64	46°45'	127°40'	12	0	—	—	—	—
6637-6648	11/11/64	46°45'	128°24'	12	3	6638	2/15/65	46°59'	124°08'
						6641	2/9/65	47°03'	124°09'
						6648	2/22/65	44°17'	124°05'
6649-6660	11/11/64	46°45'	129°36'	12	7	6649	4/14/65	48°40'	124°49'
						6651	1/14/65	49°23'	126°28'
						6652	1/23/65	48°32'	122°58'
						6656	7/31/65	51°31'	128°06'
						6657	1/15/65	49°11'	125°58'
						6658	1/10/65	49°10'	126°00'
						6659	12/29/67	49°09'	126°01'
6661-6672	11/13/64	46°45'	130°33'	12	0	—	—	—	—
6673-6684	11/14/64	47°20'	129°57'	12	1	6682	2/12/65	52°45'	129°06'
6685-6696	11/14/64	47°53'	129°15'	12	2	6685	6/28/65	53°10'	129°55'
						6687	5/22/65	52°25'	128°50'
6697-6708	11/14/64	48°10'	128°33'	12	0	—	—	—	—
6709-6720	11/15/64	48°27'	128°09'	12	1	6710	6/19/67	56°00'	133°55'
6721-6732	11/15/64	48°43'	127°37'	12	0	—	—	—	—
6733-6744	11/15/64	49°00'	127°11'	12	0	—	—	—	—
6745-6756	11/15/64	49°06'	127°00'	12	0	—	—	—	—
6757-6768	11/16/64	49°11'	126°54'	12	0	—	—	—	—
6769-6780	11/16/64	49°14'	126°46'	12	1	6770	4/5/65	52°32'	129°02'
6781-6792	11/16/64	49°20'	126°37'	12	1	6784	6/15/65	52°38'	129°20'

This study shows the seasonal reversal of flow in nearshore surface water to the geographical limits of the data, but the northerly extent of the reversal is not known at this time. The study also suggests the seaward extent of the coastal flow to be 150 to 200 km (81 to 108 nautical miles).

ACKNOWLEDGMENTS

The assistance of F. Favorite, Oceanography Program Leader, Division of Marine Fish and Shellfish, Northwest Fisheries Center, is gratefully acknowledged. The coopera-

tion of the scientific party and crew of the RV *George B. Kelez* is greatly appreciated.

LITERATURE CITED

- BENNETT, E. B.
1959. Some oceanographic features of the northeast Pacific Ocean during August 1955. *J. Fish. Res. Board Can.* 16:565-633.
- BUDINGER, T. F., L. K. COACHMAN, and C. A. BARNES.
1964. Columbia River effluent in the northeast Pacific Ocean, 1961, 1962: Selected aspects of physical oceanography. Univ. Wash., Dep. Oceanogr., Tech. Rep. 99, 78 p.

Table 4. — Release and recovery locations for drift bottles released in January 1965.

Serial number	Date released	Location		Number released	Number recovered	Serial number	Date recovered	Location	
		Lat. N	Long. W					Lat. N	Long. W
6793-6804	1/13/65	46°45'	124°18'	12	3	6793	3/11/65	51°58'	128°27'
						6796	5/3/65	48°56'	125°32'
						6797	3/6/65	—	— *
6805-6816	1/13/65	46°45'	124°36'	12	2	6811	5/17/65	46°57'	124°08'
						6814	7/11/65	49°06'	125°52'
6817-6828	1/13/65	46°45'	124°47'	12	3	6818	5/30/65	46°03'	123°57'
						6822	5/29/65	47°05'	124°08'
						6827	5/16/65	46°46'	124°04'
6829-6840	1/14/65	46°45'	124°59'	12	8	6830	1/25/65	47°41'	124°29'
						6831	1/25/65	47°41'	124°29'
						6832	1/27/65	47°37'	124°21'
						6833	1/25/65	47°41'	124°29'
						6836	1/27/65	47°54'	124°38'
						6837	1/25/65	47°40'	124°21'
						6838	1/30/65	47°40'	124°23'
						6839	1/25/65	47°41'	124°29'
6841-6852	1/14/65	46°45'	125°10'	12	5	6845	1/30/65	48°22'	124°37'
						6846	1/29/65	48°22'	124°37'
						6849	2/20/65	48°23'	124°43'
						6850	2/25/65	48°07'	123°12'
						6852	1/28/65	48°22'	124°37'
6853-6864	1/14/65	46°45'	125°20'	12	1	6854	2/15/65	46°53'	124°06'
6865-6876	1/15/65	46°45'	126°00'	12	6	6865	2/16/65	46°28'	124°03'
						6866	2/13/65	48°22'	124°37'
						6869	2/8/65	48°22'	124°37'
						6870	4/7/65	48°22'	124°38'
						6874	3/5/65	48°02'	122°37'
						6876	2/8/65	48°22'	124°37'
6877-6888	1/15/65	46°45'	126°33'	12	2	6878	4/10/65	48°42'	124°58'
						6888	2/28/65	48°22'	124°37'
6889-6900	1/15/65	46°45'	127°11'	12	4	6892	5/24/65	43°51'	124°08'
						6895	5/16/65	47°04'	124°08'
						6896	2/28/65	47°44'	124°25'
						6897	8/8/65	48°40'	124°50'
6901-6912	1/16/65	46°45'	128°24'	12	5	6903	5/15/65	46°20'	124°03'
						6904	3/26/65	46°34'	124°03'
						6905	5/2/65	46°49'	124°04'
						6906	6/27/65	49°23'	126°32'
						6911	2/27/65	47°43'	124°25'
6913-6924	1/16/65	46°45'	129°36'	12	8	6915	5/2/65	43°37'	124°10'
						6916	3/27/65	43°49'	123°57'
						6918	5/8/65	48°55'	125°31'
						6919	2/27/65	47°10'	124°09'
						6920	3/26/65	45°13'	123°56'
						6921	2/26/65	46°02'	123°53'
						6922	2/27/65	47°03'	124°08'
						6924	5/9/65	44°39'	124°03'
						6925-6936	1/16/65	46°45'	130°33'
6927	3/26/65	45°03'	123°58'						
6929	3/27/65	45°03'	123°58'						
6932	5/7/65	43°46'	124°09'						
6933	5/2/65	47°31'	124°21'						
6936	4/22/65	45°07'	123°57'						

Table 4.—Release and recovery locations for drift bottles released in January 1965, continued.

Serial number	Date released	Location		Number released	Number recovered	Serial number	Date recovered	Location	
		Lat. N	Long. W					Lat. N	Long. W
6937-6948	1/17/65	47°20'	129°57'	12	6	6937	5/17/65	46°53'	124°06'
						6938	12/7/67	49°23'	126°32'
						6939	3/17/65	46°15'	123°53'
						6940	5/17/65	45°37'	123°56'
						6941	5/9/65	46°21'	124°03'
						6944	5/14/65	47°13'	124°11'
6949-6960	1/17/65	47°53'	129°02'	12	6	6949	6/4/65	46°37'	123°58'
						6950	3/16/65	48°22'	124°37'
						6955	5/26/65	46°21'	124°02'
						6957	5/3/65	48°57'	125°35'
						6958	11/10/65	48°22'	124°37'
						6959	5/22/65	47°15'	124°11'
6961-6972	1/17/65	48°27'	128°09'	12	0	—	—	—	—
6973-6984	1/18/65	48°43'	127°37'	12	1	6977	2/8/65	50°47'	128°26'
6985-6996	1/18/65	49°00'	127°12'	12	2	6992	9/3/67	50°44'	128°23'
						6995	2/3/65	50°47'	128°26'
6997-7008	1/18/65	49°06'	127°00'	12	1	6997	8/17/65	52°11'	128°30'
7009-7020	1/18/65	49°10'	126°56'	12	0	—	—	—	—
7021-7032	1/18/65	49°14'	126°46'	12	2	7022	3/12/65	50°48'	128°26'
						7028	3/1/65	52°10'	128°30'
7033-7044	1/18/65	49°20'	126°37'	12	5	7035	7/26/65	50°01'	127°23'
						7036	5/5/65	50°03'	127°25'
						7039	1/21/65	50°00'	127°18'
						7041	3/5/65	50°03'	127°25'
						7042	9/22/65	49°58'	127°15'

*Card returned—unable to ascertain recovery location.

BURT, W. V., and B. WYATT.

1964. Drift bottle observations of the Davidson Current off Oregon. In K. Yoshida (editor), *Studies on oceanography*, p. 156-165. Univ. Tokyo Press, Tokyo.

DODIMEAD, A. J., and H. J. HOLLISTER.

1958. Progress report of drift bottle releases in the northeast Pacific Ocean. *J. Fish. Res. Board Can.* 15:851-865.

DOE, L. A. E.

1955. Offshore waters of the Canadian Pacific coast. *J. Fish. Res. Board Can.* 12:1-34.

FAVORITE, F., and D. M. FISK.

1971. Drift bottle experiments in the North Pacific Ocean and Bering Sea—1957-60, 1962, 1966, and 1970. U.S. Dep. Commer., NOAA, Natl. Mar. Fish. Serv., Data Rep. 67, 20 p. on 1 microfiche.

FAVORITE, F., D. FISK, and W. J. INGRAHAM, JR.

1965. First transponding oceanographic buoys in the Pacific. *J. Fish. Res. Board Can.* 22:689-694.

FISK, D. M.

1971. Recoveries from 1964 through 1968 of drift bottles released from a merchant vessel, S.S. *Java Mail*, enroute Seattle to Yokohama, October 1964. *Pac. Sci.* 25:171-177.

INGRAHAM, W. J., JR.

1967. The geostrophic circulation and distribution of water properties off the coasts of Vancouver Island and Washington, spring and fall 1963. U.S. Fish Wildl. Serv., Fish. Bull. 66:223-250.

REID, J. L., JR.

1960. Oceanography of the northeastern Pacific Ocean during the last ten years. *Calif. Coop. Oceanic Fish. Invest. Rep.* 7:77-90.

SCHWARTZLOSE, R. A.

1963. Nearshore currents of the western United States and Baja California as measured by drift bottles. *Calif. Coop. Oceanic Fish. Invest. Rep.* 9:15-22.

648. Weight loss of pond-raised channel catfish (*Ictalurus punctatus*) during holding in processing plant vats. By Donald C. Greenland and Robert L. Gill. December 1971, iii + 7 p., 3 figs., 2 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
649. Distribution of forage of skipjack tuna (*Euthynnus pelamis*) in the eastern tropical Pacific. By Maurice Blackburn and Michael Laurs. January 1972, iii + 16 p., 7 figs., 3 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
650. Effects of some antioxidants and EDTA on the development of rancidity in Spanish mackerel (*Scomberomorus maculatus*) during frozen storage. By Robert N. Farragut. February 1972, iv + 12 p., 6 figs., 12 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
651. The effect of premortem stress, holding temperatures, and freezing on the biochemistry and quality of skipjack tuna. By Ladell Crawford. April 1972, iii + 23 p., 3 figs., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
652. The use of electricity in conjunction with a 12.5-meter (Headrope) Gulf-of-Mexico shrimp trawl in Lake Michigan. By James E. Ellis. March 1972, iv + 10 p., 11 figs., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
653. An electric detector system for recovering internally tagged menhaden, genus *Brevoortia*. By R. O. Parker, Jr. February 1972, iii + 7 p., 3 figs., 1 appendix table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
654. Immobilization of finerling salmon and trout by decompression. By Doyle F. Sutherland. March 1972, iii + 7 p., 3 figs., 2 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
655. The calico scallop, *Argopecten gibbus*. By Donald M. Allen and T. J. Costello. May 1972, iii + 19 p., 9 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
656. Making fish protein concentrates by enzymatic hydrolysis. A status report on research and some processes and products studied by NMFS. By Malcolm B. Hale. November 1972, v + 32 p., 15 figs., 17 tables, 1 appendix table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
657. List of fishes of Alaska and adjacent waters with a guide to some of their literature. By Jay C. Quast and Elizabeth L. Hall. July 1972, iv + 47 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
658. The Southeast Fisheries Center bionumeric code. Part I: Fishes. By Harvey R. Bullis, Jr., Richard B. Roe, and Judith C. Gatlin. July 1972, xl + 95 p., 2 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
659. A freshwater fish electro-motivator (FFEM)-its characteristics and operation. By James E. Ellis and Charles C. Hoopes. November 1972, iii + 11 p., 9 figs.
660. A review of the literature on the development of skipjack tuna fisheries in the central and western Pacific Ocean. By Frank J. Hester and Tamio Otsu. January 1973, iii + 13 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
661. Seasonal distribution of tunas and billfishes in the Atlantic. By John P. Wise and Charles W. Davis. January 1973, iv + 24 p., 13 figs., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
662. Fish larvae collected from the northeastern Pacific Ocean and Puget Sound during April and May 1967. By Kenneth D. Waldron. December 1972, iii + 16 p., 2 figs., 1 table, 4 appendix tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
663. Tagging and tag-recovery experiments with Atlantic menhaden, *Brevoortia tyrannus*. By Richard L. Kroger and Robert L. Dryfoos. December 1972, iv + 11 p., 4 figs., 12 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
664. Larval fish survey of Humboldt Bay, California. By Maxwell B. Eldridge and Charles F. Bryan. December 1972, iii + 8 p., 8 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
665. Distribution and relative abundance of fishes in Newport River, North Carolina. By William R. Turner and George N. Johnson. September 1973, iv + 23 p., 1 fig., 13 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
666. An analysis of the commercial lobster (*Homarus americanus*) fishery along the coast of Maine, August 1966 through December 1970. By James C. Thomas. June 1973, v + 57 p., 18 figs., 11 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
667. An annotated bibliography of the cunner, *Tautoglabrus adspersus* (Walbaum). By Fredric M. Serchuk and David W. Frame. May 1973, ii + 43 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
668. Subpoint prediction for direct readout meteorological satellites. By L. E. Eber. August 1973, iii + 7 p., 2 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
669. Unharvested fishes in the U.S. commercial fishery of western Lake Erie in 1969. By Harry D. Van Meter. July 1973, iii + 11 p., 6 figs., 6 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
670. Coastal upwelling indices, west coast of North America, 1946-71. By Andrew Bakun. June 1973, iv + 103 p., 6 figs., 3 tables, 45 appendix figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
671. Seasonal occurrence of young Gulf menhaden and other fishes in a northwestern Florida estuary. By Marlin E. Tagatz and E. Peter H. Wilkins. August 1973, iii + 14 p., 1 fig., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
672. Abundance and distribution of inshore benthic fauna off southwestern Long Island, N.Y. By Frank W. Steimle, Jr. and Richard B. Stone. December 1973, iii + 50 p., 2 figs., 5 appendix tables.
673. Lake Erie bottom trawl explorations, 1962-66. By Edgar W. Bowman. January 1974, iv + 21 p., 9 figs., 1 table, 7 appendix tables.
674. Proceedings of the International Billfish Symposium, Kailua-Kona, Hawaii, 9-12 August 1972. Part 2. Review and Contributed Papers. Richard S. Shomura and Francis Williams (editors). July 1974, iv + 335 p., 38 papers. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
675. Price spreads and cost analyses for finfish and shellfish products at different marketing levels. By Erwin S. Penn. March 1974, vi + 74 p., 15 figs., 12 tables, 12 appendix figures, 41 appendix tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
676. Abundance of benthic macroinvertebrates in natural and altered estuarine areas. By Gill Gilmore and Lee Trent. April 1974, iii + 13 p., 11 figs., 3 tables, 2 appendix tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
677. Distribution, abundance, and growth of juvenile sockeye salmon, *Oncorhynchus nerka*, and associated species in the Naknek River system, 1961-64. By Robert J. Ellis. September 1974, v + 53 p., 27 figs., 26 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
678. Kinds and abundance of zooplankton collected by the USCG icebreaker *Glacier* in the eastern Chukchi Sea, September-October 1970. By Bruce L. Wing. August 1974, iv + 18 p., 14 figs., 6 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
679. Pelagic amphipod crustaceans from the southeastern Bering Sea, June 1971. By Gerald A. Sanger. July 1974, iii + 8 p., 3 figs., 3 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
680. Physiological response of the cunner, *Tautoglabrus adspersus*, to cadmium. October 1974, iv + 33 p., 6 papers, various authors. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
681. Heat exchange between ocean and atmosphere in the eastern North Pacific for 1961-71. By N. E. Clark, L. Eber, R. M. Laurs, J. A. Renner, and J. F. T. Saur. December 1974, iii + 108 p., 2 figs., 1 table, 5 plates.



UNITED STATES
DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
SCIENTIFIC PUBLICATIONS STAFF
ROOM 450
1107 N.E. 45TH ST.
SEATTLE, WA 98105
OFFICIAL BUSINESS

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF COMMERCE
COM-210

THIRD CLASS
BULK RATE



Marine Biological Laboratory S
Library - Periodicals
Woods Hole, Ma 02543

